



The Migratory Properties and Numbers of T Regulatory Cell Subsets in Circulation Are Differentially Influenced by Season and Are Associated With Vitamin D Status

Abigail A. Lamikanra^{1,2*}, Hoi Pat Tsang^{1,2}, Shaza Elsiddig^{1,2}, Michael Spencer^{1,2}, Elinor Curnow³, Robert Danby^{4,5} and David J. Roberts^{1,2*}

OPEN ACCESS

Edited by:

Joanna Davies,
San Diego Biomedical Research
Institute, United States

Reviewed by:

Marcin Moniuszko,
Medical University of Białystok,
Poland
Nirupama Darshan Verma,
University of New South Wales,
Australia

*Correspondence:

Abigail A. Lamikanra
abigail.lamikanra@ndcls.ox.ac.uk
David J. Roberts
david.roberts@ndcls.ox.ac.uk

Specialty section:

This article was submitted to
Immunological Tolerance
and Regulation,
a section of the journal
Frontiers in Immunology

Received: 27 September 2019

Accepted: 26 March 2020

Published: 19 May 2020

Citation:

Lamikanra AA, Tsang HP,
Elsiddig S, Spencer M, Curnow E,
Danby R and Roberts DJ (2020) The
Migratory Properties and Numbers
of T Regulatory Cell Subsets
in Circulation Are Differentially
Influenced by Season and Are
Associated With Vitamin D Status.
Front. Immunol. 11:685.
doi: 10.3389/fimmu.2020.00685

¹ National Health Service Blood and Transplant, Oxford, United Kingdom, ² Nuffield Division of Clinical Laboratory Sciences, Radcliffe Department of Medicine, University of Oxford, Oxford, United Kingdom, ³ NHS Blood and Transplant, Statistics and Clinical Studies, Bristol, United Kingdom, ⁴ Department of Haematology, Churchill Hospital, Oxford, United Kingdom, ⁵ Anthony Nolan Research Institute, London, United Kingdom

The control of peripheral immune responses by FOXP3⁺ T regulatory (Treg) cells is essential for immune tolerance. However, at any given time, Treg frequencies in whole blood can vary more than fivefold between individuals. An understanding of factors that influence Treg numbers and migration within and between individuals would be a powerful tool for cellular therapies that utilize the immunomodulatory properties of Tregs to control pathology associated with inflammation. We sought to understand how season could influence Treg numbers and phenotype by monitoring the proportion of natural thymus-derived Tregs (nTregs) defined as (CD3⁺CD4⁺CD25⁺FOXP3⁺CD127⁻/_{low}) cells as a proportion of CD4⁺ T cells and compared these to all FOXP3⁺ Tregs (allTregs, CD3⁺CD25⁺FOXP3⁺CD127⁻/_{low}). We were able to determine changes within individuals during 1 year suggesting an influence of season on nTreg frequencies. We found that, between individuals at any given time, nTreg/CD4⁺ T cells ranged from 1.8% in February to 8.8% in the summer where median nTreg/CD4 in January and February was 2.4% (range 3.75–1.76) and in July and August was 4.5% (range 8.81–3.17) $p = 0.025$. Importantly we were able to monitor individual nTreg frequencies throughout the year in donors that started the year with high or low nTregs. Some nTreg variation could be attributed to vitamin D status where normal linear regression estimated that an absolute increase in nTreg/CD4⁺ by 0.11% could be expected with 10 nmol increase in serum 25 (OH) vitamin D3 ($p = 0.005$, 95% CI: 0.03–0.19). We assessed migration markers on Tregs for the skin and/or gut. Here cutaneous lymphocyte associated antigen (CLA⁺) expression on CD25⁺FOXP3⁺CD4⁺/CD4⁺ was compared with the same population expressing the gut associated integrin, $\beta 7$. Gut tropic CD25⁺FOXP3⁺ $\beta 7$ ⁺Tregs/CD4⁺ had similar dynamics to nTreg/CD4⁺. Conversely, CD25⁺FOXP3⁺CLA⁺Tregs/CD4⁺ showed no

association with vitamin D status. Important for cellular therapies requiring isolation of Tregs, the absolute number of $\beta 7^{+}CD4^{+}CD25^{+}FOXP3^{+}$ Tregs was positively associated with 25(OH)vitamin D3 ($R^2 = 0.0208$, $r = 0.184$, $p = 0.021$) whereas the absolute numbers of $CLA^{+}CD4^{+}CD25^{+}FOXP3^{+}$ Tregs in the periphery were not influenced by vitamin D status. These baseline observations provide new opportunities to utilize seasonal variables that influence Treg numbers and their migratory potential in patients or donors.

Keywords: Tregs, regulatory T cells, vitamin D3, seasons, tolerance, migration

INTRODUCTION

Immune responses show considerable variation due to inherited and environmental factors, including not only acquired immune responses after exposure to specific antigens but also differences in individual innate responses and regulatory functions (1).

There are major patterns of development of these immune responses and functions over the lifetime of the individual, but there are also large and significant inter-individual and temporal variations (1–4). Although in principle, genetic and/or environmental factors must modulate the immune phenotype, a detailed description and quantitative understanding of the major factors involved in determining the inter-individual and temporal variations is far from complete [reviewed in ref. (5)].

$CD4^{+}CD25^{hi}$ and $FoxP3^{hi}$ T regulatory cells (Tregs) profoundly modulate effector responses, reviewed in ref. (6). In hematopoietic stem cell transplantation (HSCT) the proportion of Tregs/ $CD4^{+}$ T cells (or Tregs/ $CD3^{+}$ T cells) has a major effect on outcomes of allogeneic HSCT in animal models (7–9) and in observational studies in HSCT cohorts (10–12). In these studies, recipients of grafts with a higher than median proportion of Tregs had reduced incidence of acute graft versus host disease (aGvHD) (11) and non-relapse mortality, commonly due to complications associated with transplantation (12). Indeed, meta-analysis of observational outcomes from clinical cohorts shows that higher than median Treg/ $CD3^{+}$ T cell ratios in donor grafts are associated with reduced aGvHD, reduced transplantation-related mortality and improved overall survival (13). These initial clinical studies did not evaluate the role of naïve or memory Tregs, nor the role of Tregs that express skin-homing receptors in the graft, for example, cutaneous lymphocyte associated antigen (CLA) or gut-homing receptors such as $\alpha 4\beta 7$ integrin, on HSCT outcomes. Furthermore, the factors that determine high Treg/ $CD4^{+}$ or high Treg/ $CD3^{+}$ T cell ratios or the number and proportions of naïve, memory or tissue-specific sub-sets of Tregs in donors and in donor grafts have not been described.

There is a wide individual variation in Treg numbers and Treg/ $CD4^{+}$ T cell ratios. For example, clinical studies have shown that Treg/ $CD4^{+}$ T cell ratios vary from 0.8 to 8.6% in donor HSCT grafts (12) and similarly the Treg/ $CD4^{+}$ T cells in peripheral blood can vary from 0.2 to 7% (14–16). There are also seasonal cycles of T effector cells associated with relapse of multiple sclerosis in the summer months (17) and reviewed in ref. (18) and significant differences of innate invariant natural

killer T (i-NKT) cells that are associated with favorable outcomes following HSCT (19).

Some differences in Tregs are due to a seasonal variation in Treg/ $CD4^{+}$ ratios and expression of FoxP3 protein in humans which peak in the summer months and early autumn (June–September in the Northern Hemisphere) (2, 3, 6, 20, 21). These seasonal cycles in Treg numbers appear to follow the well-described high-summer peaks of vitamin D [see ref. (22) for review] and increased expression of the vitamin D receptor (23). Although 1, 25 dihydroxy-vitamin D may perpetuate the survival and activity of induced Tregs *in vitro* (24, 25), the studies of seasonal variation of vitamin D and Tregs have not resolved the association of vitamin D and Tregs in individuals.

Additional factors that vary with season and may also contribute to immune cell composition include urban particulate matter (PM) (26, 27) and high ambient temperature as evidenced in murine studies (28) and human studies (29). Cortisol, which forms part of the hypothalamic-pituitary-adrenal axis, is also sensitive to changes in daylight hours as well as the circadian clock, showing peaks in the first awakening hour and declining until the end of day (30–32).

We established a cohort of volunteers who were phenotyped regularly for Tregs and other immune cell types through the year to study the seasonal variation of immune cells, the numbers and phenotypes of Tregs, Treg/ $CD4^{+}$ ratios and the relationship of these immune-cell parameters with vitamin D3.

MATERIALS AND METHODS

All reagents were purchased from Sigma-Aldrich (United Kingdom) and antibodies raised in mice purchased from Becton Dickinson (BD, United Kingdom) unless stated otherwise.

Recruitment and Sample Collection

The research protocol was approved by the National Research Ethics Committee (REC reference 15/NS/0060) and the National Health Service Blood and Transplant (NHSBT) Blood Supply CARE committee. All participants gave written informed consent in accordance with the Declaration of Helsinki. A proportion of whole blood from the sampling pouch was collected into K₂EDTA vacutainer tubes (BD, Oxford, United Kingdom) and used to measure whole blood cell parameters (Sysmex XE-2100, Sysmex UK) and lymphocyte subsets and also to provide

plasma. Samples were also saved for serum using vacutainer tubes for coagulating whole blood (BD). Whole blood parameters and the frequency of Tregs, Treg subsets and i-NKT cells were measured on the same day as sample collection. Plasma was collected from the upper layer following centrifugation of whole blood for 5 min at $1000 \times g$ at room temperature, and then stored at -40°C . Serum aliquots were collected from supernatants following coagulation for 30 min in the dark at room temperature, centrifuged for 15 min at $1000 \times g$, stored at -40°C and then -80°C for long term storage after 4 months. To avoid confounding variations due to gender all data presented is from male donors.

Immune Cell Phenotyping

Whole blood was lysed using red cell lysis buffer (15 mM Ammonium Chloride, 10 mM Sodium bicarbonate, 1 mM EDTA) with gentle agitation at room temperature for 10 min. Cells were washed 3 times in PBS by centrifuging at $500 \times g$ for 5 min at 4°C , resuspended in PBS containing 0.5% BSA. Tregs were stained as follows: the cell surface phenotype was determined using an antibody cocktail that comprised BUV395 conjugated CD45 (clone H130), PE conjugated anti-human CD25 (clone M-A251), FITC conjugated anti-human CD3 (clone UCHT1), PerCpCy5.5 conjugated anti-human CD4 (clone RPA-T4) and PE-Cy7 conjugated anti CD127 (clone HIL-7R-M21). In some cocktails anti-CD127 was replaced with PE-Cy7 conjugated to anti-human CD45RA (clone HI100) or PE-Cy7 conjugated to anti-human CLA (clone HECA452) or biotinylated anti-human integrin $\beta 7$ (clone FIB504, Biolegend) followed by 1 wash and incubation with PE-Cy7 conjugated to streptavidin (eBioscience). Cells were stained for 30 min at 4°C followed by wash in ice cold 5% BSA in PBS. Finally all samples were fixed and permeabilized in the cold with the FOXP3 staining buffer set (Miltenyi Biotech) before staining for FOXP3 using APC conjugated anti-human FOXP3 (clone PCH101) from eBioscience. NK T Cells and i-NKT cells were stained with a cocktail comprising APC conjugated anti-human CD56, PE conjugated anti-human $\text{V}\alpha 24$, FITC conjugated anti-human CD3 (clone UCHT1) PerCpCy5.5 conjugated anti-human CD4 (clone RPA-T4). Before use, all antibodies were titrated for optimal working dilutions. CST bead calibration was used to normalize output from the flow cytometer to rule out batch effects due to changes in the performance of the flow cytometer. Samples incubated with the different cocktails were acquired in parallel with compensation controls and fluorescence minus one (FMO) controls for each fluorochrome using the LSR-II flow cytometer (BD). The gating used for analysis of Tregs, Treg subsets, NK and iNKT cells is outlined in **Supplementary Figure 1**. Absolute cell numbers of populations were determined using Truecount beads (BD) according to the manufacturer's guidelines.

Analysis of T Cells and Their Sub-Sets

All T cells were defined from viable DAPI negative events in the lymphocyte gate using the FSC vs. SSC plot, expression of CD45 and of CD3 (see **Supplementary Figure 1**). CD4^{+} T cells were defined from the same gate and CD8^{+} T cells estimated by subtracting the number of CD4^{+} events in the

CD3^{+} gate from the total number of CD3^{+} events in the lymphocyte gate. Thymic derived nTregs were defined as $\text{CD45}^{+}\text{CD3}^{+}\text{CD4}^{+}\text{CD25}^{+}\text{FOXP3}^{+}\text{CD127}^{-/\text{low}}$, referred to as $\text{CD4}^{+}\text{CD25}^{+}\text{FOXP3}^{+}\text{CD127}^{-/\text{low}}$ or nTreg. The mostly naïve population of Tregs were captured using $\text{CD45}^{+}\text{CD3}^{+}\text{CD4}^{+}\text{CD25}^{+}\text{FOXP3}^{+}\text{CD45RA}^{+}$ expression and referred to as $\text{CD45RA}^{+}\text{Treg}$. The proportion of skin trophic and gut trophic Tregs were defined using the same T cell markers used for naïve Tregs and replacing the antibody specific to CD45RA with antibodies to CLA and $\beta 7$, respectively. Restrictions in antibody availability, fluorochrome usage and cell numbers meant staining for the $\beta 7$ integrin was the best option to estimate $\alpha 4\beta 7$ gut associated Tregs since the frequency of all $\beta 7^{+}$ Tregs was proportional to the $\alpha 4\beta 7$ Tregs (**Supplementary Figure 2**).

Measurement of Vitamin D Metabolites and Cortisol

Serum samples were thawed to room temperature in the dark. 25-hydroxyvitamin D [$25(\text{OH})\text{D}$] was measured in duplicate using an enzyme immune assay (EIA) and following the manufacturer's guidance instructions (Immunodiagnosics, Tyne and Wear, United Kingdom). Cortisol was also measured in sera collected from the same donors and measured by EIA in duplicate following the manufacturer's (R&D Systems) instructions. For each donor and analyte, measurements from different seasons were carried out on the same plate (i.e., samples from January and February versus those from June and July) with reference samples to control for plate variation.

Acquisition of Environmental Data

Data for PM_{2.5} (particulate matter $< 2.5 \mu\text{m}$ in diameter) and PM₁₀ (particulate matter $< 10 \mu\text{m}$ in diameter) was collected from Air Quality England¹ and UK Air at the Department of the Environment, Fisheries and Rural Affairs, London: <https://ukair.defra.gov.uk/data>, accessed November 20, 2019. PM_{2.5} data for 2016 was not available in urban background or roadside sites in Oxford although PM₁₀ data was available. Therefore PM_{2.5} and PM₁₀ data reported here originates from urban stations in Oxford and the next closest station to the Oxford donor center in Reading that records this data. Data for temperature (mean monthly data) and total sunshine duration in Oxford (Location: 450900E 207200N, Lat 51.761 Lon -1.262 , 63 m above mean sea level) were obtained from the meteorological office².

Statistical Analysis

GraphPad Prism version 7 (GraphPad Software, Inc., United States) and SAS/STAT version 9.4 (SAS Institute Inc., Cary, North Carolina, United States) was used to compare the mean variables in immune cells through one-way ANOVA with Tukey's correction for multiple comparisons. Spearman's two-tailed test (abbreviated as r) was used to assess associations between dependent and independent variables. The Mann-Whitney U test was used to compare ranks within each group.

¹https://www.airqualityengland.co.uk/local-authority/?la_id=267

²<https://www.metoffice.gov.uk/pub/data/weather/uk/climate/stationdata/oxforddata.txt>

Unless otherwise stated median values with inter-quartile range (IQR) are reported in the text. Comparisons with a p -value of < 0.05 are reported as significant.

To explore whether winter and summer levels of nTreg were correlated, normal linear regression was used to describe the association between nTreg in January/February (winter) and July/August (summer). Results were adjusted for age at donation (**Supplementary Figure 3**). Mean nTreg was used for any donor with more than one nTreg measurement between January or February and December.

Normal linear regression was also used to describe the association between nTregs (in any month) and vitamin D3 level. Again, results were adjusted for age at donation. A random effect was included in this model to allow for correlation between repeat samples from the same donor. The mean of two duplicate measures of vitamin D for each donor sample was used in the analysis. All analyses were restricted to male platelet donors that were not taking vitamin D supplementation at the time of the study.

RESULTS

Characteristics of Donors

We initially recruited and consented 164 platelet donors that were healthy, eligible to donate and attended the platelet apheresis clinic between October 2015 and January 2016. Of these donors, a proportion regularly attended clinic in the morning. Samples were collected from 58 of those consecutive donors who attended clinic between January 2016 and December 2016, Monday to Friday inclusive with appointments before midday (1200). Collecting samples before midday each day facilitated the processing of samples on the same day and reduced the impact of diurnal changes on the variation of circulating immune cell numbers or characteristics (33). **Figure 1** describes the process of recruitment and selection.

We initially followed 58 donors, 7 of whom were female. As so few samples could be collected from women, due to the bias of donors eligible for donating platelets, we only analyzed samples from male donors. We were able to resample 51 Caucasian male donors, 91% of whom were resampled at least three times during the year. Three of these donors were taking supplements that contained vitamin D and were subsequently removed from the dataset. So, we had samples from 48 donors to analyse for changes in immune cell numbers and phenotypes between each month of the year. Forty-three of these donors attended the clinic in different seasons so that their individual variation in immune cell frequencies could be assessed between seasons. Mean number of visits, 5 (95% confidence interval 5–6). Samples from 32 of these donors were available to determine serum levels of vitamin D metabolites and cortisol. **Figure 1** summarizes how the samples from each participant were used.

Monthly Variation of Immune Cells Over One Year

Assessment of the absolute numbers of blood cells using the Sysmex analyzer did not reveal any seasonal change in the


white blood cell count of neutrophils, monocytes or absolute numbers of lymphocytes (**Figure 2**). Similarly, the monthly changes in absolute numbers of all T cells, CD4⁺ and CD8 T cell determined by flow cytometry were not significant (**Figures 3A–C**). However, when the numbers of thymic-derived natural Tregs (CD4⁺CD25⁺FOXP3⁺CD127^{−/low}) (nTregs) were assessed there was a gradual increase from January through to June, July and August that doubled the number of nTregs in circulation from a median of 37.4 nTregs/ μ l (IQR 71.0–34.9) to 77.3 nTregs/ μ l (IQR, 84.1–63.8), 79.2nTregs/ μ l (IQR, 90.3–61.0) and 75.2nTregs/ μ l (IQR, 91.7–61.8), $p < 0.01$. The decline in nTreg numbers from July and August to December (median of 56.5 nTregs/ μ l, IQR 70.8–41.0) was also apparent ($p < 0.03$) (**Figure 3D**).

We next assessed the FOXP3⁺ Tregs within both the CD4⁺ and CD8⁺ compartment (CD3⁺CD25⁺FOXP3⁺CD127^{−/low}) and described as all Tregs from now on. We compared all Tregs/CD3⁺ T cells with nTreg/CD4⁺ T cells and with CD4⁺V α 24⁺/CD4⁺ T cells to represent a subset of immune-regulatory i-NKT cells.

The median proportion of all Treg cells/CD3⁺ T cells increased substantially from 2.5% (IQR, 2.9–1.7) in January and February to then almost double to 4.9% (IQR 6.0–4.0) in July and August ($p < 0.001$) after which they declined from September and October when they were 4.7% of CD3⁺ T cells (IQR 5.3–3.9), to 3.7% (IQR 4.1–3.2) in November and December ($p = 0.004$) (**Figure 4A**). Similarly, the proportion of nTreg/CD4⁺ T cells also increased from a median of approximately 3.4% (IQR 3.9–2.7) at the start of the year to a median of 4.0% (IQR 4.6–3.4) in July and August ($p = 0.025$) to then decrease back to 3.0% (IQR 3.5–2.6) at the end of the year ($p = 0.004$) (**Figure 4B**).

The proportion of CD4⁺ T cells expressing the V α 24⁺ chain showed negligible seasonal change where median values were the same at 0.5% of all CD4⁺ T cells. A comparison of means showed that in May and June, the mean proportion of CD4⁺ T cells with the V α 24⁺ T cell chain was 0.58% and median was 0.5% (IQR 0.7–0.4). This then decreased in November and December where the mean of V α 24⁺ T cells was 0.47% but median was again 0.5% (IQR 0.6–0.4) $p = 0.027$ (**Figure 4C**).

Thus, the concentration of all FOXP3⁺ Tregs showed the greatest variation throughout the year of the typed sub-sets of peripheral blood mononuclear cells. We wanted to know if this was also true for the relative levels of nTregs in each participant as most literature refers to nTregs as *bone fide* Tregs. **Figure 4D** shows the distribution of nTregs in individuals that had annual nTreg levels below the 25th percentile (Q1), between the 25th and 50th percentile (Q2), between the 50th and 75th percentile (Q3) and above the 75th percentile (Q4). Those with average nTregs that fell below the annual median of 3.5% nTreg/CD4⁺ had nTreg frequencies that remained mostly below or at the 50th percentile throughout the year. The exception was between May and August where for those in which measurements were taken, nTreg/CD4⁺ increased by 1%. Conversely 50% of individuals with average nTregs above the 50th or 75th percentile started the year with nTreg/CD4⁺ between 4 and 4.5% that remained at or above median from April onward.



Moreover, using a linear regression model, we found that higher nTreg/CD4⁺ in winter was associated with higher nTregs in summer ($p = 0.028$). An increase in nTreg/CD4⁺ of 1% in January/February was associated with a 0.4% increase in nTreg/CD4⁺ in July/August (estimate, 95% confidence interval: 0.37%, 0.04–0.69%).

Variation of Treg Subsets Over the Year

We first looked at Treg subsets as a proportion of all CD4⁺ T cells as this analysis would allow assessment of their relative contribution to the total T helper cell population (Figure 5).

CD45RA⁺ Tregs

The CD45RA⁺ mostly naïve population of Tregs (CD3⁺CD4⁺CD25⁺FOXP3⁺CD45RA⁺) varied considerably

between individuals and within certain months of the year. Nonetheless we found that on average the median proportion of these Tregs were at their lowest in November and December (0.4% CD45RA⁺Tregs/CD4⁺), IQR 0.6–0.2 compared with March and April (0.6%, IQR 0.9–0.4), $p = 0.004$ or July and August (0.6%, IQR 0.9–0.4), $p = 0.037$. See Figure 5A.

Tregs With Tissue Specific Homing Markers

In order to capture Tregs that could migrate to epithelial mucosa in the intestines, the contribution of (CD3⁺CD4⁺CD25⁺FOXP3⁺β7⁺) β7⁺Tregs to the CD4⁺ T cell population was determined. The proportion of β7⁺Treg/CD4⁺ T cells almost doubled from winter months (0.4%, IQR 0.5–0.2) to July, August (0.7%, IQR 1–0.4) $p < 0.0001$, with noticeably higher levels persisting in September and October (0.6%, IQR




FIGURE 2 | Monthly changes in whole blood parameters determined using an automated hematological analyzer. The variation in numbers of white blood cells (WBC), lymphocytes, monocytes and neutrophils are shown.

0.9–0.5), $p = 0.002$ (Figure 5B). The median proportion of Tregs within the CD4^+ compartment that expressed the skin trophic marker CLA ($\text{CD3}^+\text{CD4}^+\text{CD25}^+\text{FOXP3}^+\text{CLA}^+$)/ CD4^+ T cells was unchanged from January and February (1.8%, IQR 2.7–1.4) until May and June (1.85%, IQR 2.5–1.4). $\text{CLA}^+\text{Treg}/\text{CD4}^+$ then declined from July and August (1.8%, IQR 2.2–1.1) through to the end of the year in November and December (1.1%, IQR 1.5–0.90), $p < 0.0001$ (Figure 5C).

We also assessed the absolute cell numbers of these subsets and observed that the median number of CD45RA^+ Treg cells ($\text{CD3}^+\text{CD4}^+\text{CD25}^+\text{FOXP3}^+\text{CD45RA}^+$)/ μL were lowest in January/February (7/ μL , IQR 12–4) and in November/December (8/ μL , IQR 12–4) with a trend to peak in July and August (11 cells/ μL IQR, 17–6), $p = 0.07$ (Figure 5D). Similarly the median number of $\beta 7^+$ Treg cells ($\text{CD3}^+\text{CD4}^+\text{CD25}^+\text{FOXP3}^+\beta 7^+$)/ μL increased from January and February (9 cells/ μL IQR, 12–6) to July and August (13 cells/ μL IQR, 16–8), $p = 0.006$ to then decreased in November and December to 8 cells/ μL , IQR, 11–6), $p = 0.001$ (Figure 5E). In contrast the median number of CLA^+ Treg cells ($\text{CD3}^+\text{CD4}^+\text{CD25}^+\text{FOXP3}^+\text{CLA}^+$)/ μL were three to four fold greater in number than CD45RA^+ or $\beta 7^+$ Tregs/ μL and did not vary from January and February (at 39 cells/ μL IQR, 48–29) to July and August (39 cells/ μL IQR, 50–27). However, there was a significantly reduced number of the CLA^+ subset by November and December (27 CLA^+ Treg cells/ μL , IQR 36–20, $p = 0.0004$).

Treg Subsets as a Proportion of Total Tregs

There was no obvious change in the proportion of naïve CD45RA^+ Tregs/($\text{CD3}^+\text{CD4}^+\text{CD25}^+\text{FOXP3}^+$)Tregs, median values of which were between 14.5% and 15.4% ($p = 0.9$) (Figure 6A). Although there was a trend to increase $\beta 7^+$ Tregs/Tregs between May/June (12.6%, IQR 16.4–7.9) and September and October (17%, IQR 20.25–13.9), this was not significant, $p = 0.60$ (Figure 6B). Conversely the median proportion of skin trophic CLA^+ Tregs/Tregs decreased by almost 15% between January/February (59.1%, IQR 66.1–54.9) through to September/October (42.8%, IQR 51.6–36.5), $p < 0.0001$ (Figure 6C).

We assessed the same Treg subsets in each of the donors with multiple measurements in the same year. The mean annual value of Treg subset for each donor was used to assign each person to quartiles as described for nTregs in Figure 4D. Overall, individuals with CD45RA^+ Tregs/Tregs and $\beta 7^+$ Tregs/Tregs in the lowest two quartiles (Q1 and Q2) were more likely to remain relatively low throughout the year (Figures 6D,E) as were those with CLA^+ Tregs/Tregs (Figure 6F).

Association of Seasonal Variables With Tregs

Vitamin D

To determine which factors may influence the observed differences in nTreg and Treg subsets we examined the role




FIGURE 3 | Variation in absolute numbers of T cells each month. The estimate of absolute numbers of T cell subsets are shown for each month and were determined after immune-phenotyping and analysis as described in Materials and Methods for all CD3⁺ T cells (**A**), CD4⁺ T cells (**B**), CD8⁺ T cells (**C**), and nTregs (**D**). In **D** the *p*-values are shown using one way ANOVA analysis of log transformed values where the *p*-values for changes in nTregs/ μ L between January and June, July and August are 0.010, 0.004, and 0.004, respectively; between July and December *p* = 0.032 and between August and December, *p* = 0.026.


of endogenous vitamin D (**Figure 7**). Solar UVB irradiation mediates the conversion of 7-dehydrocholesterol to VD3 in the skin which is then converted to 25(OH)VD3 in the liver or in peripheral myeloid cells. Given that the number of daylight hours in the same year followed a similar distribution to the proportion of FOXP3⁺ Tregs (**Figures 7A, 3, 4**) we measured 25(OH) vitamin D metabolite in consecutive samples from donors that attended clinic and observed gradual increases from January through to September (**Figure 7B**). 25(OH) vitamin D was positively associated with all Tregs ($R^2 = 0.165$, $r = 0.406$, $p < 0.0001$) and with thymic-derived nTregs ($R^2 = 0.042$, $r = 0.178$, $p = 0.023$) (**Figures 7C,D**).

Conversely, there was no positive association between 25(OH) vitamin D and CD45RA⁺Treg/CD4⁺ ($R^2 = 0.008$, $r = 0.085$, $p = 0.284$) nor CLA⁺Treg/CD4⁺ ($R^2 = 0.0011$, $r = 0.051$, $p = 0.519$). See **Figures 7E,G**. However, $\beta 7^+$ Treg/CD4⁺ were positively associated with vitamin D status ($R^2 = 0.067$, $r = 0.267$, $p = 0.001$) again relating seasonal changes of this subset of

Tregs with observed changes in vitamin D status of this population (**Figure 7F**).

Data was adjusted for age and repeated measurements in the same donor as described in methods, and an association between nTreg/CD4⁺ and vitamin D ($p = 0.0052$) observed. An increase in vitamin D of 10 nmol was associated with an absolute 0.11% increase in nTreg/CD4⁺ T cell percentage (estimate, 95% confidence interval: 0.11, 0.03–0.19).

In keeping with the observation that the proportion of Tregs that are either CD45RA⁺ or $\beta 7^+$ appear mostly constant throughout the year, we observed no association of CD45RA⁺Tregs/Tregs ($R^2 = 0.008$, $r = 0.069$, $p = 0.388$) nor of $\beta 7^+$ Treg/Tregs ($R^2 = 0.002$, $r = 0.057$, $p = 0.476$) (**Figures 8A,B**) with 25 (OH) vitamin D3. In contrast, the CLA⁺ fraction of Tregs CLA⁺Tregs/Tregs was negatively associated with 25(OH) vitamin D in sera ($R^2 = 0.029$, $r = -0.167$, $p = 0.034$) (**Figure 8C**). Conversely the absolute numbers of CLA⁺ Tregs (CLA⁺Treg cells/ μ L) did not change according to vitamin D status, indicating



that the number of Tregs expressing CLA were not influenced by vitamin D ($R^2 = 0.010$, $r = 0.056$, $p = 0.488$) (**Figure 8F**). However, the absolute numbers of $\beta 7^+$ Tregs were positively associated with vitamin D status ($R^2 = 0.021$, $r = 0.184$, $p = 0.021$) as were CD45RA⁺Tregs ($R^2 = 0.021$, $r = 0.152$, $p = 0.052$) (**Figures 8D,E**).

Cortisol


Since endogenous cortisol levels are influenced by sunlight hours, we also examined the association of total cortisol in each participant with the frequency of Tregs and their subsets. Cortisol was measured in samples from the same cohort that was used to measure 25(OH) vitamin D. We observed that the lowest levels of cortisol were in the summer months (May through to August) and highest levels in November and December (**Figure 9A**). Subsequently Cortisol in this cohort showed weak but significantly negative correlations with all FOXP3⁺ T cells/CD3 ($R^2 = 0.041$, $r = -0.222$, $p = 0.005$) and with thymic-derived nTregs/CD4⁺ ($R^2 = 0.061$, $r = -0.257$, $p = 0.001$) (**Figures 9B,C**). Similarly CD45RA⁺Treg/CD4⁺ and $\beta 7^+$ Treg/CD4⁺ were negatively associated with cortisol in sera ($R^2 = 0.055$, $r = -0.320$, $p < 0.0001$ and $R^2 = 0.018$, $r = -0.221$, $p = 0.005$, respectively) but CLA⁺Treg/CD4⁺ ($R^2 = 0.009$, $r = -0.132$, $p = 0.094$) were not (**Figures 9D–F**). With the exception of CD45RA⁺Treg/Treg ($R^2 = 0.016$, $r = -0.189$, $p = 0.016$) we observed no correlation between

subsets as a proportion of Tregs (**Figures 10A–C**). However, the absolute numbers of CD45RA⁺ Treg and $\beta 7^+$ Treg cells/ μ L ($R^2 = 0.020$, $r = -0.171$, $p = 0.030$ and $R^2 = 0.023$, $r = -0.165$, $p = 0.039$, respectively) were weakly but significantly associated with cortisol (**Figures 10D,E**) whereas there was no correlation of CLA⁺Treg cells/ μ L ($R^2 = 0.005$, $r = -0.023$, $p = 0.771$) with cortisol (**Figure 10F**).

DISCUSSION

Previous reports have described inter-individual variation of immune cells providing a rich resource to understand the influence of environment and genetic traits on different branches of immunity (3, 4). On the whole, the variation in T cell immunity and the cytokines that support their variability is largely governed by non-inheritable traits.

Here, we examined both inter-individual and intra-individual variation of FOXP3⁺ immuno-regulatory T cells and their subsets with changes in season in male platelet donors that returned to clinic on average five times a year. We found that both absolute numbers and the proportion of nTreg/CD4⁺ T cells were higher in the summer than in the winter. This was also true for allTreg/CD3 and was consistent with changes in vitamin D status and the ability of vitamin D to induce FOXP3 expression (24). In contrast similar changes to CD4⁺T cells expressing the




Vα24⁺ chain (i-NKT/CD4⁺) were almost undetectable. Our data showing variation of nTreg within each donor from one season to the next allowed us to estimate that an increase of 10 nM 25 (OH) vitamin D3 in serum could result in a 0.1% absolute increase of nTreg as a percentage of CD4⁺ T cells. In this study we were able to monitor the intra-individual variation of Tregs to reduce any influence of genetic factors on our observations.

We also described changes in the CD45RA⁺ subset of FOXP3⁺ Tregs that can define the stability of Treg function. CD45RA has been used to isolate Tregs that retain immunomodulatory activity after expansion *in vitro* implying longer lived suppression *in vivo* (34, 35). We found that CD45RA⁺ Tregs/CD4⁺ remained at their highest levels between Spring and Autumn and represented less than 1% of the T-helper population at their peak. However, the proportion of CD45RA⁺ Tregs within the FOXP3⁺ Treg population did not vary, so supporting previous observations that the apparent decrease in CD45RA⁺ Tregs/CD4⁺ T from Autumn to Winter is due to the increased proliferative potential of CD45RA-memory Tregs rather than a change in output from the thymus (36).

Examination of the potential of Tregs to migrate to gut mucosa (β7⁺ Tregs) or to the skin (CLA⁺ Tregs) suggests that the dynamics of these two populations are affected differently by seasonal variables. The proportion of β7⁺ Tregs within the T helper population resembled the month-to-month dynamics of nTreg/CD4⁺ and was positively associated with vitamin D status. Conversely, the dynamics CLA⁺ Tregs/CD4⁺ differed considerably; between January and June CLA⁺ Tregs/CD4⁺ and were on average twice the levels observed in November and December.

When we examined the composition of Tregs we found that the CLA⁺ Tregs/Tregs appeared to vary, whilst the proportion of β7⁺ and CD45RA⁺ within Tregs were mostly unchanged. This could in part be explained by the observed increase in absolute numbers of Tregs expressing the β7 integrin between January and July/August thereby reducing the proportion of CLA⁺ cells in the Treg population which were unchanged during the same period (Figures 5E,F). The absolute numbers of Tregs expressing CLA did not change in association with vitamin D3 in




sera whilst the number of Tregs expressing β 7 integrin or CD45RA did change.

A recent study has shown that β 7⁺ Tregs are CD45RA⁺ and comprise new emigrants from the thymus where expression of β 7 is induced by IL-2 (37). CD45RA⁺ Tregs are considered most suitable for adoptive cellular therapy given that they retain suppressive function upon expansion (34, 38, 39). It is possible that the increase in β 7⁺ Treg subsets is in part due to conversion of naïve β 7⁺ CD4⁺ T cells to β 7⁺FOXP3⁺ Tregs following increased exposure to sunlight and induced expression of VDR or similarly induced from the activated memory population of β 7⁺CD4⁺ T cells (40).

We were unable to confirm CD45RA⁺ expression on CD3⁺CD4⁺CD25⁺FOXP3⁺CD127^{low} β 7⁺ Tregs

(CD45RA⁺ β 7⁺ Tregs) nor assess the effect of season on CD45RA⁺ β 7⁺ Tregs in our cohort. However, analysis of this population in freshly isolated PBMCs from 6 donors shows that CD45RA⁺ β 7⁺ Tregs are on average 1/5th of β 7⁺ Tregs (Supplementary Figure 4) indicating that the majority of β 7⁺Tregs are likely effector memory Tregs and would be less stable if expanded *ex vivo*.

Brodin et al. (1) have also reported that the frequency of naïve T cells and their response to IL-2 is one of the few strongly heritable factors that can influence an immune response. The apparently positive but weak association of CD45RA⁺ numbers with vitamin D3 may also be dependent on the responsiveness of Tregs to IL-2 within each participant of this study. Nonetheless future application of season and vitamin




D3 to enhance $\beta 7^+$ Tregs will be dependent on establishing the stability of suppressive function of any of the induced Treg populations.

The variation in expression of CLA within the FOXP3⁺CD4⁺CD25⁺Treg population was considerable, both between and within individuals. The gradual intra-individual decrease from January through to mid-summer and November of the proportion of CLA⁺ Tregs may be due to migration of CLA⁺ Tregs from the periphery to the skin in response to increased ultraviolet (UV) exposure from the sun (36, 41). This could again partly explain the marked decrease in CLA⁺Tregs/total Tregs in blood samples collected between March and September when exposure to sunlight had increased.

The poor recovery of CLA⁺ Tregs in samples collected when daylight hours were reduced in autumn and winter may be due to a delay in migration back to the periphery and also the decline in all Tregs since 1, 25 (OH) vitamin D can induce or stabilize FOXP3 (24) (U. Leuschner, personal communication) and with fewer daylight hours the potential to do so would be reduced.

Interestingly, a study of 350 older male and female participants aged between 69 and 80, has shown that the fraction of CLA⁺Tregs in circulating Tregs is positively associated with the degree of skin pigmentation following exposure to UV radiation from the sun in Florida (42). In this location daylight hours were between 800 and 500 hours from July 2014 to July 2015. This was at least double the number of hours in Oxford from January 2016 to December 2016 which were between 400 and 200 daylight hours.³ A correction for sex was applied to this data but data from male and female volunteers was not analyzed separately. Furthermore, the authors did not report vitamin D metabolites nor did they report exclusion of data from volunteers that were taking vitamin D supplements. Thus, whilst the data from this study provides compelling evidence for the ability of UV exposure to influence the migration of Tregs the conclusions presented therein are unlikely to apply to all demographics.

³<http://www.timebie.com/sun> and <https://www.timeanddate.com/sun> accessed December 2018.



Other studies have also reported changes in Treg frequencies and phenotype with season using flow cytometry (4, 20, 21). None of these studies observed an association between Treg frequencies and vitamin D status nor did they relate vitamin D3 to Tregs within the same person during the course of 1 year.


The possible explanations for the different observations presented here include differences in the cohorts studied. Our data is from volunteers with an older mean age of 55 (median 58, range 21 to 72). Several studies have shown that age can influence the number and activation of different T cells especially baseline levels of Tregs and their naïve status [see Brodin et al for review (5)] (3, 4, 43). Also, we restricted our analysis to data from male volunteers to remove any confounding effects due to imbalance of genders since differences in hormones and the menstrual cycle can influence the relative proportion of Tregs in circulation (44) reviewed by (5). Variation of diurnal effects on Tregs (23) were reduced by collecting all samples between 8 am and 12 noon and measuring Treg numbers within 2 h of collection. Importantly, we defined the Treg population in our samples by selecting CD25⁺FOXP3⁺CD127^{low} cells which would influence the specificity of the associations that we observed compared to studies that used different markers to stain for Tregs. Finally we were able to minimize the influence of genetic traits and age by periodically sampling from the same donors so that we could determine

the association between nTreg and 25 (OH) vitamin D in each donor during 1 year.

A number of studies that utilize Mass Cytometry to define different Treg populations suggest that there can be overlap between Tregs and different Thelper cell subsets (45, 46) and these would not have been identified with the antibodies used in this study. Our study did not differentiate between effector memory and central memory Tregs. The use of additional phenotype markers to characterize the memory compartment in more detail using CD45RO, CD62L and specific chemokine receptor markers would have provided a more detailed understanding of the change in dynamics within the Treg memory and effector memory population.

Our inability to detect changes in the proportions of iNKT cells may have been due to the very low frequency of these cells in circulation and the reduced sensitivity of the approach we used to detect them which would require dual staining for V β 11 TCR chain (47). Due to ethical considerations we were unable to measure the Treg levels and phenotype in the skin of healthy volunteers in this study to better explain the decrease in CLA⁺Tregs in peripheral blood.

We examined the association of vitamin D status on Treg frequencies as this would be the most tractable method to enhance Tregs in HSCT donors. However, environmental factors, other than vitamin D status and exposure to UV, are




likely to contribute to the seasonal variability observed. Due to ethical considerations we were unable to retrospectively determine the exposure of individual participants to sun. Questionnaires to address this should be employed in future studies. External factors that could influence Treg frequencies include ambient temperature which shows the same pattern as sunlight hours and vitamin D status and is therefore also positively associated with Treg frequencies (see **Supplementary Figure 5**). Conversely PM_{2.5} and PM₁₀ may negatively affect Treg levels (**Supplementary Figure 6**). In order to associate these observations more accurately with the Treg frequency of each donor the ambient temperature around the time of sample collection and PM levels local to each participant should also be measured.

We were however able to relate cortisol levels to Treg frequencies within participants that attended more than twice in 1 year. Here we observed negative associations with the frequencies of all Tregs and nTregs as well as absolute numbers of CD45RA⁺ and $\beta 7^+$ Tregs. This could explain the marked

reduction of Tregs in November and December when cortisol levels were at their highest in this cohort.

There is evidence to support the interaction between the hypothalamus-pituitary axis, vitamin D₃ and T cell frequencies. A high density of VDR and vitamin D₃ metabolizing enzymes are present within the hypothalamus (48). Salivary cortisol can be reduced by 40% in healthy male and female volunteers following vitamin D₃ supplementation with 2000 IU/day ($n = 9$) versus placebo ($n = 6$) for 14 days (49) and by 15% following supplementation of with 4,000 IU/day ($n = 22$) versus placebo ($n = 19$) for 16 weeks in relapsing and remitting patients with multiple sclerosis (50). However, these responses are likely to be most relevant during the first hour of awakening since the dip in cortisol over the course of 1 day remained unchanged between treatment and placebo (50).

Pre-clinical experimental models indicate that cortisol decreases Tregs through binding to the mineral corticoid receptor which enhances pro-inflammatory mediators [see ref. (51) for review]. Furthermore, the proportion of Tregs and



their expression of FOXP3 is decreased in healthy volunteers whilst T helper 17 (Th17) cells are enhanced immediately after completing endurance exercise; and cortisol in serum is increased alongside IL-6, IL-10, and TGF- β . IL-6 and TGF- β are required to generate pathogenic Th17 cells and IL-6 in the sera of endurance-athletes may contribute to the reduced expansion of Tregs *in vitro* (52). These observations suggest that cortisol inhibits Tregs indirectly through enhancement of inflammatory mediators and/or support of Th17 expansion. Since the participants in our study were not taking vitamin D supplements, it is possible that the higher levels of cortisol compounded the low vitamin D status in November and December to reduce Treg numbers through enhancement of pro-inflammatory mediators.

Experimental models of Treg phenotype and expansion *in vivo* have shown that metabolites of vitamin A and short chain fatty acids can alter the migratory properties and numbers of Tregs reviewed in refs. (53) and (54, 55). In humans, exposure to allergens, different infections and the gut microbiome influence Treg numbers (3, 5) and seasonal variations in melatonin may influence Treg survival by inhibiting Th17 T cell development (17). The investigation of these possibilities in future studies

would improve our understanding of how the balance of T regulatory cells can be modified for application in the clinic.

Treg numbers in HSCT grafts may vary from 0.8% nTreg/CD4 to 8.6% nTreg/CD4 and nTreg/CD4 frequencies above 3% can double overall survival (11, 12). Our observation that a 0.11% increase in nTreg requires a 10 nmol/L increase in VitD3 level indicates that to increase nTreg:CD4 from 2 to 3% (an increase of 1%) would require an increase of ≥ 91 nmol/L (i.e., $[1/0.11 \times 10]$) 25(OH) vitamin D3 in HSCT donors with Treg < 2%. A placebo controlled trial (56) suggests that a one-off dose of 250,000 IU vitamin D3 could safely increase 25 (OH) vitamin D3 by this amount (from 41.5 nmol/L to 103.2 nmol/L (CI 87.1–122.2 nmol/L) for 5–10 days during the winter months). This relationship may be modified in female donors and it will be important to establish this to better understand how to use vitamin D3 to induce Treg subsets.

In this study we have found that enhanced vitamin D status in healthy males was associated with changes in the frequency of different FOXP3⁺ Treg populations in the periphery that could be modified by levels of cortisol. An increase in the frequency of nTregs and $\beta 7$ ⁺Tregs

in the CD4⁺ T helper population and of FOXP3⁺ Tregs in the T cell population was more apparent in the summer months than in the winter months. The opposite was true of the proportion of Tregs in the periphery that expressed CLA as these were not associated with the vitamin D status of the cohort studied here. Since an increase in the frequency of circulating Tregs in the recipient that are CLA⁺ or $\alpha 4\beta 7^{+}$ is associated with reduced risk of skin or gut GvHD, respectively (57, 58) the baseline data presented here will support endeavors to improve clinical outcomes following HSCT. Further investigations into how appropriate supplementation with vitamin D3 could modify cortisol levels, the migratory potential and suppressive stability of Tregs in HSCT are therefore warranted.

DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding author.

ETHICS STATEMENT

The studies involving human participants were reviewed and approved by North of Scotland Research Ethics Service (NHS Grampian). The participants provided their written informed consent to take part in this study.

REFERENCES

1. Brodin P, Jojic V, Gao T, Bhattacharya S, Angel CJ, Furman D, et al. Variation in the human immune system is largely driven by non-heritable influences. *Cell*. (2015) 160:37–47. doi: 10.1016/j.cell.2014.12.020
2. Tsang JS, Schwartzberg PL, Kotliarov Y, Biancotto A, Xie Z, Germain RN, et al. Global analyses of human immune variation reveal baseline predictors of postvaccination responses. *Cell*. (2014) 157:499–513. doi: 10.1016/j.cell.2014.03.031
3. Carr EJ, Dooley J, Garcia-Perez JE, Lagou V, Lee JC, Wouters C, et al. The cellular composition of the human immune system is shaped by age and cohabitation. *Nat Immunol*. (2016) 17:461–8. doi: 10.1038/ni.3371
4. Aguirre-Gamboa R, Joosten I, Urbano PCM, van der Molen RG, van Rijssen E, van Cranenbroek B. Differential effects of environmental and genetic factors on T and B cell immune traits. *Cell Rep*. (2016) 17:2474–87. doi: 10.1016/j.celrep.2016.10.053
5. Brodin P, Davis MM. Human immune system variation. *Nat Rev Immunol*. (2017) 17:21–9. doi: 10.1038/nri.2016.125
6. Sakaguchi S, Miyara M, Costantino CM, Hafler DA. FOXP3⁺ regulatory T cells in the human immune system. *Nat Rev Immunol*. (2010) 10:490–500. doi: 10.1038/nri2785
7. Taylor PA, Lees CJ, Blazar BR. The infusion of ex vivo activated and expanded CD4⁺CD25⁺ immune regulatory cells inhibits graft-versus-host disease lethality. *Blood*. (2002) 99:3493–9.
8. Hoffmann P, Ermann J, Edinger M, Fathman CG, Strober S. Donor-type CD4⁺CD25⁺ regulatory T cells suppress lethal acute graft-versus-host disease after allogeneic bone marrow transplantation. *J Exp Med*. (2002) 196:389–99.
9. Edinger M, Hoffmann P, Ermann J, Drago K, Fathman CG, Strober S, et al. CD4⁺CD25⁺ regulatory T cells preserve graft-versus-tumor activity while inhibiting graft-versus-host disease after bone marrow transplantation. *Nat Med*. (2003) 9:1144–50.

AUTHOR CONTRIBUTIONS

AL designed and implemented the study, analyzed data, prepared figures, and wrote the manuscript. HT performed the experiments and analyzed the data. SE and MS performed the experiments. EC performed the statistical analysis of data. RD contributed to the design of the study. DR conceived, designed the study, and wrote the manuscript.

FUNDING

This work was supported by the NHS Blood and Transplant intramural R&D funding.

ACKNOWLEDGMENTS

We would like to thank Drs. John Girdlestone, Nick Ciccone, and Letizia Lo Faro for their critical review during the preparation of this manuscript and Mrs. Wendy Slack for administrative support and proof reading.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2020.00685/full#supplementary-material>

10. Di Ianni M, Falzetti F, Carotti A, Terenzi A, Castellino F, Bonifacio E, et al. Tregs prevent GVHD and promote immune reconstitution in HLA-haploidentical transplantation. *Blood*. (2011) 117:3921–8. doi: 10.1182/blood-2010-10-311894
11. Pastore D, Delia M, Mestice A, Carluccio P, Perrone T, Gaudio F, et al. CD3⁺/Tregs ratio in donor grafts is linked to acute graft-versus-host disease and immunologic recovery after allogeneic peripheral blood stem cell transplantation. *Biol Blood Marrow Transplant*. (2012) 18:887–93. doi: 10.1016/j.bbmt.2011.10.039
12. Danby RD, Zhang W, Medd P, Littlewood TJ, Peniket A, Rocha V, et al. High proportions of regulatory T cells in PBSC grafts predict improved survival after allogeneic haematopoietic SCT. *Bone Marrow Transplant*. (2016) 51:110–8. doi: 10.1038/bmt.2015.215
13. Fisher SA, Lamikanra A, Doree C, Gratton B, Tsang P, Danby RD, et al. Increased regulatory T cell graft content is associated with improved outcome in haematopoietic stem cell transplantation: a systematic review. *Br J Haematol*. (2017) 176:448–63. doi: 10.1111/bjh.14433
14. Zhang W, Smythe J, Frith E, Belfield H, Clarke S, Watt SM, et al. An innovative method to generate a good manufacturing practice-ready regulatory T-cell product from non-mobilized leukapheresis donors. *Cytotherapy*. (2015) 17:1268–79. doi: 10.1016/j.jcyt.2015.05.015
15. Blache C, Chauvin JM, Marie-Cardine A, Contentin N, Pommier P, Dedreux I, et al. Reduced frequency of regulatory T cells in peripheral blood stem cell compared to bone marrow transplantations. *Biol Blood Marrow Transplant*. (2010) 16:430–4. doi: 10.1016/j.bbmt.2009.10.027
16. Gregg R, Smith CM, Clark FJ, Dunnion D, Khan N, Chakraverty R, et al. The number of human peripheral blood CD4⁺ CD25^{high} regulatory T cells increases with age. *Clin Exp Immunol*. (2005) 140:540–6.
17. Farez MF, Mascanfroni ID, Mendez-Huergo SP, Yeste A, Murugaiyan G, Garo LP, et al. Melatonin contributes to the seasonality of multiple sclerosis relapses. *Cell*. (2015) 162:1338–52. doi: 10.1016/j.cell.2015.08.025

18. Watad A, Azrielant S, Soriano A, Bracco D, Abu Much A, Amital H. Association between seasonal factors and multiple sclerosis. *Eur J Epidemiol.* (2016) 31:1081–9.
19. Chaidos A, Patterson S, Szydio R, Chaudhry MS, Dazzi F, Kanfer E, et al. Graft invariant natural killer T-cell dose predicts risk of acute graft-versus-host disease in allogeneic hematopoietic stem cell transplantation. *Blood.* (2012) 119:5030–6. doi: 10.1182/blood-2011-11-389304
20. Schweintzger NA, Gruber-Wackernagel A, Shirsath N, Quehenberger F, Obermayer-Pietsch B, Wolf P. Influence of the season on vitamin D levels and regulatory T cells in patients with polymorphic light eruption. *Photochem Photobiol Sci.* (2016) 15:440–6. doi: 10.1039/c5pp00398a
21. Khoo AL, Koenen HJ, Chai LY, Sweep FC, Netea MG, van der Ven AJ, et al. Seasonal variation in vitamin D(3) levels is paralleled by changes in the peripheral blood human T cell compartment. *PLoS One.* (2012) 7:e29250. doi: 10.1371/journal.pone.0029250
22. Wacker M, Holick MF. Sunlight and Vitamin D: a global perspective for health. *Dermatoendocrinol.* (2013) 5:51–108. doi: 10.4161/derm.24494
23. Dopico XC, Evangelou M, Ferreira RC, Guo H, Pekalski ML, Smyth DJ, et al. Widespread seasonal gene expression reveals annual differences in human immunity and physiology. *Nat Commun.* (2015) 6:7000. doi: 10.1038/ncomms8000
24. Kang SW, Kim SH, Lee N, Lee WW, Hwang KA, Shin MS, et al. 1,25-Dihydroxyvitamin D3 promotes FOXP3 expression via binding to vitamin D response elements in its conserved noncoding sequence region. *J Immunol.* (2012) 188:5276–82. doi: 10.4049/jimmunol.1101211
25. Unger WW, Laban S, Kleijwegt FS, van der Slik AR, Roep BO. Induction of Treg by monocyte-derived DC modulated by vitamin D3 or dexamethasone: differential role for PD-L1. *Eur J Immunol.* (2009) 39:3147–59. doi: 10.1002/eji.200839103
26. Pfeffer PE, Ho TR, Mann EH, Kelly FJ, Sehlstedt M, Pourazar J, et al. Urban particulate matter stimulation of human dendritic cells enhances priming of naive CD8 T lymphocytes. *Immunology.* (2018) 153:502–12. doi: 10.1111/imm.12852
27. Sarkar S, Rivas-Santiago CE, Ibironke OA, Carranza C, Meng Q, Osornio-Vargas A, et al. Season and size of urban particulate matter differentially affect cytotoxicity and human immune responses to *Mycobacterium tuberculosis*. *PLoS One.* (2019) 14:e0219122. doi: 10.1371/journal.pone.0219122
28. Moriyama M, Ichinohe T. High ambient temperature dampens adaptive immune responses to influenza A virus infection. *Proc Natl Acad Sci USA.* (2019) 116:3118–25. doi: 10.1073/pnas.1815029116
29. Lee MK, Xu CJ, Carnes MU, Nichols CE, Ward JM, The Bios Consortium, et al. Genome-wide DNA methylation and long-term ambient air pollution exposure in Korean adults. *Clin Epigenetics.* (2019) 11:37. doi: 10.1186/s13148-019-0635-z
30. Hadlow NC, Brown S, Wardrop R, Henley D. The effects of season, daylight saving and time of sunrise on serum cortisol in a large population. *Chronobiol Int.* (2014) 31:243–51. doi: 10.3109/07420528.2013.844162
31. Persson R, Garde AH, Hansen AM, Osterberg K, Larsson B, Orbaek P, et al. Seasonal variation in human salivary cortisol concentration. *Chronobiol Int.* (2008) 25:923–37. doi: 10.1080/0742052080253648
32. Walker BR, Best R, Noon JP, Watt GC, Webb DJ. Seasonal variation in glucocorticoid activity in healthy men. *J Clin Endocrinol Metab.* (1997) 82:4015–9.
33. Druzd D, Matveeva O, Ince L, Harrison U, He W, Schmal C, et al. Lymphocyte circadian clocks control lymph node trafficking and adaptive immune responses. *Immunity.* (2017) 46:120–32. doi: 10.1016/j.immuni.2016.12.011
34. Hoffmann P, Eder R, Boeld TJ, Doser K, Pishesha B, Andreesen R, et al. Only the CD45RA+ subpopulation of CD4+CD25high T cells gives rise to homogeneous regulatory T-cell lines upon in vitro expansion. *Blood.* (2006) 108:4260–7.
35. Miyara M, Yoshioka Y, Kitoh A, Shima T, Wing K, Niwa A, et al. Functional delineation and differentiation dynamics of human CD4+ T cells expressing the FoxP3 transcription factor. *Immunity.* (2009) 30:899–911. doi: 10.1016/j.immuni.2009.03.019
36. Booth NJ, McQuaid AJ, Sobande T, Kissane S, Agius E, Jackson SE, et al. Different proliferative potential and migratory characteristics of human CD4+ regulatory T cells that express either CD45RA or CD45RO. *J Immunol.* (2010) 184:4317–26. doi: 10.4049/jimmunol.0903781
37. Hsu S, Lai CL, Hu M, Santner-Nanan B, Dahlstrom JE, Lee CH, et al. IL-2 enhances gut homing potential of human naive regulatory T cells early in life. *J Immunol.* (2018) 200:3970–80. doi: 10.4049/jimmunol.1701533
38. Canavan JB, Scotta C, Vossenkamper A, Goldberg R, Elder MJ, Shoval I, et al. Developing in vitro expanded CD45RA+ regulatory T cells as an adoptive cell therapy for Crohn's disease. *Gut.* (2016) 65:584–94. doi: 10.1136/gutjnl-2014-306919
39. Arroyo Hornero R, Betts GJ, Sawitzki B, Vogt K, Harden PN, Wood KJ. CD45RA distinguishes CD4+CD25+CD127-/low TSDR demethylated regulatory T cell subpopulations with differential stability and susceptibility to tacrolimus-mediated inhibition of suppression. *Transplantation.* (2017) 101:302–9. doi: 10.1097/TP.0000000000001278
40. Kongsbak M, von Essen MR, Boding L, Levring TB, Schjerling P, Lauritsen JP, et al. Vitamin D up-regulates the vitamin D receptor by protecting it from proteasomal degradation in human CD4+ T cells. *PLoS One.* (2014) 9:e96695. doi: 10.1371/journal.pone.0096695
41. Omland SH, Nielsen PS, Gjerdrum LM, Gniadecki R. Immunosuppressive environment in basal cell carcinoma: the role of regulatory T cells. *Acta Derm Venereol.* (2016) 96:917–21. doi: 10.2340/00015555-2440
42. Hesterberg RS, Amorrtu RP, Zhao Y, Hampras S, Akuffo AA, Fenske N, et al. Subpopulations associated with recent ultraviolet radiation exposure in a skin cancer screening Cohort. *J Immunol.* (2018) 201:3269–81. doi: 10.4049/jimmunol.1800940
43. Seddiki N, Santner-Nanan B, Tangye SG, Alexander SI, Solomon M, Lee S, et al. Persistence of naive CD45RA+ regulatory T cells in adult life. *Blood.* (2006) 107:2830–8.
44. Arruvito L, Sanz M, Banham AH, Fainboim L. Expansion of CD4+CD25+and FOXP3+ regulatory T cells during the follicular phase of the menstrual cycle: implications for human reproduction. *J Immunol.* (2007) 178:2572–8.
45. Kordasti S, Costantini B, Seidl T, Perez Abellan P, Martinez Llordella M, McLornan D, et al. Deep phenotyping of Tregs identifies an immune signature for idiopathic aplastic anemia and predicts response to treatment. *Blood.* (2016) 128:1193–205. doi: 10.1182/blood-2016-03-703702
46. Kunicki MA, Amaya Hernandez LC, Davis KL, Bacchetta R, Roncarolo MG. Identity and diversity of human peripheral Th and T regulatory cells defined by single-cell mass cytometry. *J Immunol.* (2018) 200:336–46. doi: 10.4049/jimmunol.1701025
47. Tocheva AS, Mansour S, Holt TG, Jones S, Chancellor A, Sanderson JP, et al. The clonal invariant NKT cell repertoire in people with Type 1 diabetes is characterized by a loss of clones expressing high-affinity TCRs. *J Immunol.* (2017) 198:1452–9. doi: 10.4049/jimmunol.1600255
48. Smolders J, Schuurman KG, van Strien ME, Melief J, Hendrickx D, Hol EM, et al. Expression of vitamin D receptor and metabolizing enzymes in multiple sclerosis-affected brain tissue. *J Neuropathol Exp Neurol.* (2013) 72:91–105. doi: 10.1097/NEN.0b013e31827f4fcc
49. Al-Dujaili EA, Munir N, Iniesta RR. Effect of vitamin D supplementation on cardiovascular disease risk factors and exercise performance in healthy participants: a randomized placebo-controlled preliminary study. *Ther Adv Endocrinol Metab.* (2016) 7:153–65. doi: 10.1177/2042018816653357
50. Rolf L, Damoiseaux J, Huitinga I, Kimenai D, van den Ouweland J, Hupperts R, et al. Stress-Axis regulation by Vitamin D3 in multiple sclerosis. *Front Neurol.* (2018) 9:263. doi: 10.3389/fneur.2018.00263
51. Belden Z, Deilius JA, Dobre M, Rajagopalan S. The role of the mineralocorticoid receptor in inflammation: focus on kidney and vasculature. *Am J Nephrol.* (2017) 46:298–314. doi: 10.1159/000480652
52. Perry C, Pick M, Bdoelach N, Hazan-Halevi I, Kay S, Berr I, et al. Endurance exercise diverts the balance between Th17 cells and regulatory T cells. *PLoS One.* (2013) 8:e74722. doi: 10.1371/journal.pone.0074722
53. Mora JR, Iwata M, von Andrian UH. Vitamin effects on the immune system: vitamins A and D take centre stage. *Nat Rev Immunol.* (2008) 8:685–98. doi: 10.1038/nri2378

54. Arpaia N, Campbell C, Fan X, Dikiy S, van der Veeken J, deRoos P, et al. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature*. (2013) 504:451–5. doi: 10.1038/nature12726
55. Smith PM, Howitt MR, Panikov N, Michaud M, Gallini CA, Bohlooly YM, et al. The microbial metabolites, short-chain fatty acids, regulate colonic Treg cell homeostasis. *Science*. (2013) 341:569–73. doi: 10.1126/science.1241165
56. Kearns MD, Binongo JN, Watson D, Alvarez JA, Lodin D, Ziegler TR, et al. The effect of a single, large bolus of vitamin D in healthy adults over the winter and following year: a randomized, double-blind, placebo-controlled trial. *Eur J Clin Nutr*. (2015) 69:193–7. doi: 10.1038/ejcn.2014.209
57. Engelhardt BG, Jagasia M, Savani BN, Bratcher NL, Greer JP, Jiang A, et al. Regulatory T cell expression of CLA or alpha(4)beta(7) and skin or gut acute GVHD outcomes. *Bone Marrow Transplant*. (2011) 46:436–42. doi: 10.1038/bmt.2010.127
58. Issa F, Robb RJ, Wood KJ. The where and when of T cell regulation in transplantation. *Trends Immunol*. (2013) 34:107–13. doi: 10.1016/j.it.2012.11.003

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Lamikanra, Tsang, Elsiddig, Spencer, Curnow, Danby and Roberts. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.